arXiv:1602.03657v1 [math.PR] 11 Feb 2016

SYMMETRIES AND MARTINGALES IN A STOCHASTIC MODEL FOR THE NAVIER-STOKES EQUATION

RÉMI LASSALLE AND ANA BELA CRUZEIRO

Abstract: A stochastic description of solutions of the Navier-Stokes equation is investigated. These solutions are represented by laws of finite dimensional semi-martingales and characterized by a weak Euler-Lagrange condition. A least action principle, related to the relative entropy, is provided. Within this stochastic framework, by assuming further symmetries, the corresponding invariances are expressed by martingales, stemming from a weak Noether's theorem.

Keywords: Stochastic analysis; Stochastic control; Navier-Stokes equation.
Mathematics Subject Classification : 93E20, 60H30

Contents

1. The weak stochastic Euler-Lagrange condition. 2
1.1. Admissible trajectories. 2
1.2. A weak description of random trajectories. 3
1.3. A weak Euler-Lagrange condition 3
2. Navier-Stokes equation and the weak Euler-Lagrange condition 3
2.1. Description of dissipative flows by laws of semi-martingales 4
2.2. Stochastic action and relative entropy 4
2.3. Least action principle and relative entropy. 4
3. Invariances and the stochastic Noether theorem. 5
3.1. Symmetry by translation and the momentum process 6
3.2. Symmetry by rotation and the kinetic momentum process 6
References 7

Several stochastic models for the *Navier-Stokes* equation have been proposed in the literature. Some refer to random perturbations of the fluid velocity. This is not the case here: we are interested in stochastic Lagrangian paths whose (mean) velocity, or drift, represent the deterministic velocity of the fluid. Different studies of stochastic Lagrangian paths in fluid dynamics and in particular in turbulence can be found in a collection of works, from which we refer to [2], [15] and [16] as examples. Also representation formulae in terms of different random processes were given in [5], [6], [8], among others.

Concerning the derivation of solutions of Navier-Stokes equations from (stochastic) variational principles, after the early articles [17] and [20], such principles were developed in [3] and subsequent works. We mention also [11] and [9] for different, unrelated approaches to the same kind of problems.

In [7] and [13] a weak description of a stochastic deformation of mechanics has been investigated: the Euler-Lagrange condition extends as a condition on laws of stochastic processes. The main originality of

this approach, with respect to other weak deformations of mechanics, is to handle problems in a functional-analytic framework, in a full consistency with the deterministic case.

In this paper, within the framework of [7] and [13], we develop a specific case, from an example of [7], which provides a *stochastic description* for solutions to the *Navier-Stokes equation*. Then we investigate several *symmetries*, whose associated invariances correspond in this setup to *martingales*.

Section 1 fixes the framework and notations of the paper; the *weak Euler-Lagrange condition* of [13] and [7] is recalled. Under conditions, in Section 2, a map

$$P : u \to P_u,$$

associates laws of $\mathbb{R}^d-$valued *semi-martingales* to *divergence free vector fields*. Solutions of the *Navier-Stokes equation* are shown to be divergence free vector fields u, whose associated probability P_u satisfies a weak *Euler-Lagrange condition* (Proposition 1); Corollary 2 characterizes those solutions as *critical points* of the *stochastic action*

$$\mathcal{S}^p(\nu) := E_\nu \left[\int_0^1 \left(\frac{|v_s^\nu|^2}{2} - p(1-t, W_t) \right) dt \right],$$

where ν denotes the *law* of specific continuous *semi-martingales*, and (v_t^ν) the characteristic drift of ν, as stated accurately below. The function p is a smooth *pressure field* which is assumed to be *given*.

The action functional above is related to the relative *entropy* with respect to a reference law μ_p induced by the pressure field.

Finally, within this stochastic model, Section 3 investigates *invariances*, stemming from *symmetries*, by the weak *Noether's theorem* of [7]; within this stochastic framework *martingales* on the canonical space play the rôle of constants of motion in classical mechanics.

1. The weak stochastic Euler-Lagrange condition.

The *weak* stochastic *Euler-Lagrange condition*, recalled below, was introduced in [7],[13]. It *embeds* in probability measures, specifically in a set of *laws* of semi-martingales, the classical condition. Thus, it provides a *functional analytic* approach to tackle *stochastic variational problems*; in particular, in contrast with usual diffusion approaches, it directly embeds the not stochastic case. In this context the extension of *Noether's theorem* becomes natural. Moreover, as stated in [7], another specificity of this framework is that it provides critical conditions to *semi-martingale optimal transportation problems*. The latter, introduced in [18], correspond to a relaxation of a specific *dynamical Schrödinger problems* (see [14]) by allowing, in particular, the characteristic dispersion to be not-trivial. Finally, as it is expected of optimization over a subset of Borel probabilities on a Polish space, one crucial advantage of this framework is that *compactness* is rather simple to obtain.

1.1. **Admissible trajectories.**

Trajectories of infinitely small passive tracers in fluids can be described by elements in the space $W := C([0,1], \mathbb{R}^d)$, namely the set of continuous $\mathbb{R}^d$-valued paths (where we consider the norm $|.|_W$ of uniform convergence), endowed with the Borel sigma-field $\mathcal{B}(W)$. In particular, trajectories of finite energy can be described by the subset

$$H := \left\{ h \in W, h := \int_0^\cdot \dot{h}_s ds, \int_0^1 |\dot{h}_s|^2_{\mathbb{R}^d} ds < \infty \right\},$$

of absolutely continuous paths with square integrable derivatives.

Let $(W_t)_{t\in[0,1]}$ denote the *evaluation process*

$$(t, \omega) \in [0,1] \times W \to W_t(\omega) := \omega(t) \in \mathbb{R}^d.$$

We consider $(\mathcal{F}_t^0)$ the natural (past) filtration and denote by $\mathcal{P}_W$ the set of Borel probabilities on W.

1.2. **A weak description of random trajectories.** In order to avoid measurability issues, we consider $(\mathcal{F}_t^\nu)$, the $\nu-$usual augmentation of the filtration $(\mathcal{F}_t^0)$ under $\nu \in \mathcal{P}_W$. The latter naturally models random trajectories, since any $\nu \in \mathcal{P}_W$ is the law of the evaluation process, on the completed probability space $(W, \mathcal{B}(W)^\nu, \nu)$. We define $\mathbb{S}$ to be the subset of $\nu \in \mathcal{P}_W$ such that there exists a $(\mathcal{F}_t^\nu)-$martingale (M_t^ν) on $(W, \mathcal{B}(W)^\nu, \nu)$, which satisfies

$$W_t = W_0 + M_t^\nu + \int_0^t v_s^\nu ds,$$

for all $t \in [0,1]$, $\nu-a.s.$, where (v_s^ν) is a $(\mathcal{F}_t^\nu)-$*predictable process* on the same space, and where the *predicable covariation process* of (M^ν) is of the specific form

$$< (M_t^\nu)^i, (M_t^\nu)^j >= \int_0^\cdot (\alpha_s^\nu)^{ij} ds,$$

for a predictable process (α_s^ν); subsequently, by abuse of language, we refer to (v_t^ν, α_t^ν) as the *characteristics* of ν. In the whole paper, notations are those of [7].

1.3. **A weak Euler-Lagrange condition.** Given a *smooth Lagrangian* function

$$\mathcal{L} : (t,x,v,a) \in [0,1] \times \mathbb{R}^d \times \mathbb{R}^d \times (\mathbb{R}^d \otimes \mathbb{R}^d) \to \mathcal{L}_t(x,v,a) \in \mathbb{R}, \tag{1.1}$$

the classical *Euler-Lagrange condition* naturally extends to $\mathbb{S}$ (see [13]). A semi-martingale $\nu \in \mathbb{S}$ satisfies the Euler-Lagrange condition if there exists a $(\mathcal{F}_t^\nu)$ *càd-làg martingale* (N_t^ν), such that

$$\partial_v \mathcal{L}_t(W_t, v_t^\nu, \alpha_t^\nu) - \int_0^t \partial_q \mathcal{L}_s(W_s, v_s^\nu, \alpha_s^\nu) ds = N_t^\nu \quad \lambda \otimes \nu - a.e., \tag{1.2}$$

$\partial_q \mathcal{L}$ and $\partial_v \mathcal{L}_t$ denoting the respective gradients (in the first and in the second variables, respectively).

Remark 1. *Similar conditions were considered in* [1] *for arbitrary semi-martingales U on abstract stochastic basis $(\Omega, \mathcal{A}, (\mathcal{A}_t), \mathcal{P})$. On the contrary, condition (1.2) imposes constraints on laws of processes. A semi-martingale U on an arbitrary stochastic basis whose law satisfies (1.2) exhibits very precise properties, depending on the Lagrangian; for instance, in a specific case, it is associated to systems of coupled stochastic differential equations; the latter are not satisfied, in general, when U verifies the critical condition of* [1]. *Moreover, the associated variational principles of* [1] *do not contain the optimum criticality for semi-martingale optimal transportation problems either.*

2. Navier-Stokes equation and the weak Euler-Lagrange condition

Henceforth, and *until the end of the paper, p denotes a smooth map*

$$p : (t,x) \in [0,1] \times \mathbb{R}^d \to p(t,x) \in \mathbb{R}^+, \tag{2.3}$$

which is further assumed to be bounded, with bounded derivatives; p models the *pressure field*. This map being given, we provide a *stochastic model* for solutions of the equation

$$\partial_t u + (u.\nabla)u = -\nabla p + \frac{\Delta u}{2} \ ; \ div \ u = 0. \tag{2.4}$$

For sake of clarity, we focus on the case where the *divergence free velocity vector field*, involved in the Navier-Stokes equation, belongs to

$$C^{1,2}_{b,div}([0,1] \times \mathbb{R}^d) := \{u \in C^{1,2}([0,1] \times \mathbb{R}^d; \mathbb{R}^d) \cap C_b([0,1] \times \mathbb{R}^d; \mathbb{R}^d) : div \ u(t,.) = 0, \ for \ all \ t \in [0,1]\}.$$

2.1. **Description of dissipative flows by laws of semi-martingales.** Given $u \in C^{1,2}_{b,div}([0,1]\times\mathbb{R}^d)$, we define P_u to be the probability measure, which is equivalent with respect to the *Wiener measure* $\mu \in \mathcal{P}_W$ (the law of *standard Brownian motion*), with density defined by

$$\frac{dP_u}{d\mu} := \exp\left(-\int_0^1 u(1-t,W_t)dW_t - \frac{1}{2}\int_0^1 |u(1-t,W_t)|^2 dt\right).$$

By the *Girsanov theorem* (see for example [10]), we obtain a map

$$P : u \in C^{1,2}_{b,div}([0,1]\times\mathbb{R}^d) \to P_u \in \mathbb{S}, \tag{2.5}$$

such that $\lambda \otimes P_u$ *a.e.*, $\alpha_s^{P_u} = I_{\mathbb{R}^d}$, and $v_t^{P_u} = -u(1-t,W_t)$, λ denoting the Lebesgue measure. *Itô's formula* yields the following:

Proposition 1. *A time-dependent vector field $u \in C^{1,2}_{b,div}([0,1]\times\mathbb{R}^d)$ satisfies the Navier-Stokes equation (2.4) if and only if P_u satisfies (1.2) for*

$$\mathcal{L}_t^p(x,v,a) := \frac{|v|^2}{2} - p(1-t,x). \tag{2.6}$$

□

2.2. **Stochastic action and relative entropy.** Define the *stochastic action* associated to the Lagrangian $\mathcal{L}^p$ of (2.6) by

$$\mathcal{S}^p : \nu \in \mathbb{S} \to \mathcal{S}^p(\nu) := E_\nu\left[\int_0^1 \mathcal{L}_s^p(W_s, v_s^\nu, \alpha_s^\nu)ds\right] \in \mathbb{R}\cup\{+\infty\}. \tag{2.7}$$

As $\mathcal{L}^p$ and P_u satisfy the assumptions of Theorem 5.1 of [7], for all p as above and $u \in C^{1,2}_{b,div}([0,1]\times\mathbb{R}^d)$, the $\mathbb{S}-$functional $\mathcal{S}^p$ is differentiable in the sense considered in [7]. Thus, we obtain the following result :

Proposition 2. *A vector field $u \in C^{1,2}_{b,div}([0,1]\times\mathbb{R}^d)$ is a solution of the equation (2.4) if and only if*

$$\delta\mathcal{S}_{P_u}[h] = 0,$$

for all $(\mathcal{F}_t^{P_u})-$adapted process (h_t) of finite energy, such that

$$h_0 = h_1 = 0 \ P_u - a.s.,$$

where $P_u \in \mathbb{S}$ is given by (2.5) and $\delta\mathcal{S}_{P_u}$ denotes the $\mathbb{S}-$differential of [7] *at P_u.* □

2.3. **Least action principle and relative entropy.** Subsequently, assuming $\nu \in \mathcal{P}_W$ to be absolutely continuous with respect to a reference law $\eta \in \mathcal{P}_W$, the relative *entropy* of ν w.r.t. η is defined as

$$\mathcal{H}(\nu|\eta) := E_\nu\left[\ln\frac{d\nu}{d\eta}\right].$$

By the representation formula in [12], we obtain

$$\mathcal{S}^p(P_u) = \mathcal{H}(P_u|\mu_p) + \ln Z_p, \tag{2.8}$$

for all $u \in C^{1,2}([0,1]\times\mathbb{R}^d)$, where P_u is defined by (2.5), and where μ_p is the absolutely continuous probability with respect to the *standard Wiener measure* ($W_0 = 0$, $\mu - a.s.$), whose density is given by

$$\frac{d\mu_p}{d\mu} := \frac{\exp\left(\int_0^1 p(1-s,W_s)ds\right)}{Z_p},$$

with Z_p a normalization constant. Whence we obtain the following ersatz of Proposition 2:

Proposition 3. *A time-dependent vector field $u \in C^{1,2}_{b,div}([0,1]\times\mathbb{R}^d)$ is a solution of the equation (2.4) if and only if*

$$\delta\mathcal{H}(.|\mu_p)_{P_u}[h] = 0$$

for all $(\mathcal{F}^\nu_t)-$adapted process (h_t) of finite energy, such that

$$h_0 = h_1 = 0 \quad P_u - a.s..$$

□

3. Invariances and the stochastic Noether theorem.

Within this section we consider the case $d = 3$ and we denote by (e_1, e_2, e_3) the canonical orthogonal basis of $\mathbb{R}^3$. We further assume that $u \in C^{1,2}_{b,div}([0,1]\times\mathbb{R}^3)$ is a solution of (2.4) for a given smooth function p. By Proposition 1, the associated law of the continuous semi-martingale P_u (c.f. (2.5)) satisfies (1.2) for $\mathcal{L}^p$ defined by (2.6).

The next subsections investigate different *symmetries* and compute the related local *martingales*, stemming from the *weak Noether Theorem* 6.1. of [7]. In each particular case considered below *the symmetries are expressed through a condition on the pressure field p*. Given the associated family of transformations (h^ϵ), subsequently, the *symmetry condition* on p yields that (h^ϵ) is a smooth family of $\mathbb{S}-$invariant transformations for $\mathcal{L}^p$, in the sense considered in [7].

We recall this *symmetry condition* on $\mathbb{S}$. First, by setting

$$\Gamma^\epsilon : \omega \in W \to \Gamma^\epsilon(\omega) \in W,$$

where

$$\Gamma^\epsilon_t(\omega) := h^\epsilon(t, \omega(t)),$$

for all $t \in [0,1]$, $\omega \in W$, (h^ϵ) induces a family (Γ^ϵ) of transformations of W. Given $\eta \in \mathbb{S}$, for all ϵ, (Γ^ϵ_t) defines a stochastic process on the probability space $(W, \mathcal{B}(W)^\eta, \eta)$. Thus, by *Itô's formula* on the probability space $(W, \mathcal{B}(W)^\eta, \eta)$, for all $\epsilon \in \mathbb{R}$, *the transformation h^ϵ of the state space $\mathbb{R}^3$ is lifted to a transformation*

$$\eta \in \mathbb{S} \to \Gamma^\epsilon_\star \eta \in \mathbb{S}$$

of $\mathbb{S}$, by pushforward. The *symmetry condition* considered in [7] consists in the relation

$$\mathcal{L}^p_t(W_t, v^\eta_t, \alpha^\eta_t) = \mathcal{L}^p_t(\Gamma^\epsilon_t, v^{\Gamma^\epsilon_\star \eta}_t \circ \Gamma^\epsilon, \alpha^{\Gamma^\epsilon_\star \eta}_t \circ \Gamma^\epsilon), \tag{3.9}$$

holding a.e., for all $\eta \in \mathbb{S}$ in the domain of the map defined in (2.7), and for all $\epsilon \in \mathbb{R}$. Here $\circ$ denotes the pullback of the ($\Gamma^\epsilon_\star\eta-$equivalence class of) map(s) $v^{\Gamma^\epsilon_\star \eta}_t : W \to \mathbb{R}^3$ with the ($\eta-$ equivalence class of) map(s) $\Gamma^\epsilon : W \to W$.

Consider a smooth Lagrangian $\mathcal{L}$ and assume that $\nu \in \mathbb{S}$ satisfies the weak Euler-Lagrange condition for this Lagrangian. The stochastic weak Noether's Theorem in [7] associates to a family (h_ϵ) of $\mathbb{S}$-invariant transformations of $\mathcal{L}$ local martingales on the probability space $(W, \mathcal{B}(W)^\nu, \nu)$. These local martingales, that we denote by $(\mathcal{I}_t)_{t\in[0,1)}$, are in fact explicitly given:

$$\mathcal{I}_t := < \frac{d}{d\epsilon}|_{\epsilon=0} h^\epsilon_t(W_t), \phi^\nu_t >_{\mathbb{R}^d} - \sum_i \left[\frac{d}{d\epsilon}|_{\epsilon=0} h^\epsilon_\cdot(W_\cdot)^i, \phi^{\nu\, i}_\cdot\right]_t + \int_0^t \theta_s ds \tag{3.10}$$

where $[.,.]$ stands for the quadratic co-variation process of *càd-làg* semi-martingales, (ϕ^ν_t) denotes a *càdlàg* modification of the process $\partial_v\mathcal{L}_t(W_t, v^\nu_t, \alpha^\nu_t)$, and

$$\theta_s := \sum_{i,j} \kappa^{i,j}_s \frac{\partial\mathcal{L}}{\partial\alpha_{i,j}}(W_s, v^\nu_s, \alpha^\nu_s), \tag{3.11}$$

where $(\kappa_s(\omega))$ is the $\mathcal{M}_d(\mathbb{R})-$valued process defined by

$$\kappa_s(\omega) := \alpha_s^{\nu}.\left(\left(\nabla\frac{d}{d\epsilon}h^{\epsilon}|_{\epsilon=0}\right)(s,W_s)\right)^{\dagger} + \left(\left(\nabla\frac{d}{d\epsilon}h^{\epsilon}|_{\epsilon=0}\right)(s,W_s)\right)\alpha_s^{\nu}. \tag{3.12}$$

3.1. **Symmetry by translation and the momentum process.** Assume the *symmetry by translation* along e_3 of the pressure, that is

$$p(t,x+ae_3) = p(t,x),$$

for all $a\in\mathbb{R}$, $t\in[0,1]$, $x\in\mathbb{R}^3$. To check that Noether's theorem yields the expected result, set

$$h^{\epsilon} : (t,x)\in[0,1]\times\mathbb{R}^3 \to h^{\epsilon}(t,x) := x+\epsilon e_3\in\mathbb{R}^3.$$

By proposition 3.2. of [7], (3.9) is trivially satisfied, so that, by Theorem 6.1 of [7], we obtain $(<v_t^{P_u},e_z>)$, as the related $(\mathcal{F}_t^{P_u})-$*local martingale*.

3.2. **Symmetry by rotation and the kinetic momentum process.** Assume the *symmetry by rotation* along the axis e_3 of the pressure; that is,

$$p(t,R^{\epsilon}x) = p(t,x), \tag{3.13}$$

for all $(t,x)\in[0,1]\times\mathbb{R}^3$ and $\epsilon\in\mathbb{R}$, where $R^{\epsilon}:\mathbb{R}^3\to\mathbb{R}^3$ denotes the operator of rotation, along the axis e_3, with angle ϵ. We consider the family of space transformations

$$h^{\epsilon} : (t,x)\in[0,1]\times\mathbb{R}^3 \to h^{\epsilon}(t,x) := R^{\epsilon}x\in\mathbb{R}^3. \tag{3.14}$$

Applying Lemma 3.2. of [7], with h^{ϵ} given by (3.14), we compute the characteristics of $\Gamma_{\star}^{\epsilon}\eta$ and we obtain the relation

$$|v^{\Gamma_{\star}^{\epsilon}\eta}\circ\Gamma^{\epsilon}|_{\mathbb{R}^3} = |v^{\eta}|_{\mathbb{R}^3} \quad \lambda\otimes\eta-a.e.,$$

for any $\epsilon\in\mathbb{R}$. Whence, the *symmetry condition* (3.9) is satisfied, from (3.13). Define the stochastic process (l_t), on the complete probability space $(W,\mathcal{B}(W)^{P_u},P_u)$, by

$$l_t := <W_t,e_1>_{\mathbb{R}^3}<v_t^{P_u},e_2>_{\mathbb{R}^3}^2 - <W_t,e_2>_{\mathbb{R}^3}<v_t^{P_u},e_1>_{\mathbb{R}^3}$$

for $t\in[0,1]$, the stochastic counterpart to the *kinetic momentum* along e_3. Further denoting by

$$rot\ u_t : \mathbb{R}^3\to\mathbb{R}^3,$$

the rotational of $u(t,.)$, the weak Noether Theorem 6.1 of [7] implies that the corresponding process $(\mathcal{I}_t)$, defined by

$$\mathcal{I}_t := l_t + \int_0^t <rot\ u_{1-s}(W_s),e_3>ds,$$

is the *local martingale* associated to this *symmetry by rotation*; otherwise stated, the stochastic kinetic momentum process l_t along e_3 is a semi-martingale on the canonical filtered probability space of ν, whose finite variation term is determined by the rotational of u along the symmetry axis. The latter expresses the *dissipation*, modeled through the martingale part of P_u, which is involved in the covariation process in the expression of $\mathcal{I}_t$.

References

[1] Bismut, J.-M. *Mécanique aléatoire*, Lecture Notes in Mathematics, 866 Springer (1981)

[2] Bossy, M., Jabir, J.F., Talay, D. *On conditional McKean Lagrangian stochastic models*, Prob. Th. and Rel. Fields 151, 319-351 (2011)

[3] Cruzeiro, A.B., Cipriano, F., *Navier-Stokes equation and diffusions on the group of homeomorphisms of the torus*, Comm. Math. Phys. 275, 255-269 (2007)

[4] Cruzeiro, A.B., Lassalle, R. *On the least action principle for the Navier-Stokes equation* Springer Proceedings in Mathematics and Statistics, Vol. 100 (2014)

[5] Constantin, P. *An Eulerian-Lagrangian approach to the Navier-Stokes equations*, Comm. Math. Phys. 216 (3), 663686 (2001)

[6] Constantin, P.Iyer, G. *A stochastic Lagrangian representation of the three-dimensional incompressible Navier-Stokes equations*, Comm. Pure Appl. Math. Phys. 61 (3), 330-345 (2008)

[7] Cruzeiro, A.B., Lassalle, R. *Weak calculus of variations for functionals of laws of semi-martingales*, submitted (2015)

[8] Cruzeiro, A.B., Shamarova, E. *Navier-Stokes equations and forward-backward SDEs on the group of diffeomorphisms of a torus*, Stoch. Proc. Appl. 119 (12), 4034-4060 (2009)

[9] Gomes, D.A., *A variational formulation for the Navier-Stokes equation*, Comm. Math. Phys. 257 (1), 227-234 (2005)

[10] Ikeda, N. and Watanabe, S., *Stochastic Differential Equations and Diffusion Processes*, North Holland, Amsterdam (Kodansha Ltd., Tokyo) (1981)

[11] Inoue, A., Funaki, T., *A new derivation of the Navier-Stokes equation*, Comm. Math. Phys. 65, 83-90 (1979)

[12] Föllmer, H. *Random fields and diffusion processes*, Ecole d' Été de Saint Flour XV–XVII (1988)

[13] Lassalle, R., Zambrini, J.C. *A weak approach to the stochastic deformation of classical mechanics* (2015)

[14] Leonard, C. *A survey of the Schrödinger problem and some of its connections with optimal transport*, Discrete and Cont. Dyn. Syst. A, 34 (4), 1533 (2014)

[15] Mikulevicius, R., Rozovskii, B. L, *Stochastic Navier-Stokes equations for turbulent flows*, SIAM J. Math. Anal. 35 (5) 1250-1310 (2004)

[16] Mikulevicius, R., Rozovskii, B.L., *On equations of stochastic fluid mechanics*, Stochastics in Finite and Infinite Dimensions, T. Hida, R. Karandikar, H. Kunita, B. Rajput, S. Watanabe, and J. Xiong, eds., Birkhauser Boston, Boston, MA, 285-302 (2001)

[17] Nakagomi, T. Yasue, K., Zambrini, J.C. *Stochastic variational derivations of the Navier-Stokes equation*, Lett. Math. Phys. 5 (6), 545-552 (1981)

[18] Tan, X., Touzi, N.*Optimal transportation under Controlled stochastic dynamics* Annals of Probability. (2012)

[19] Thieullen, M. Zambrini, J.C. *Probability and quantum symmetries I, the theorem of Noether in Schrödinger's Euclidean Quantum Mechanics*, Ann. Inst. H.Poincaré, Phys. Theo. 67 (3), 297 (1997)

[20] Yasue, K. *A variational principle for the Navier-Stokes equation*, J. Funct. Anal. 51 (2), 133-141 (1983)

[21] Zambrini, J.C. *Stochastic mechanics according to E. Schrödinger*, Phys. Rev. A. 33 (3), 1532-1548(1986)

(Rémi Lassalle) CEREMADE Université Paris-Dauphine Place du Maréchal de Lattre de Tassigny 75016 Paris

E-mail address: lassalle@ceremade.dauphine.fr

(Ana Bela Cruzeiro) GFMUL and Departamento Matemática Instituto Superior Técnico, Univ. de Lisboa, Av. Rovisco Pais, 1049-001 Lisbon, Portugal

E-mail address: abcruz@math.tecnico.ulisboa.pt